

- Tanaka, Y., 1958: Seasonal changes occurring in the gonad of *Stichopus japonicus*. — Bull. Fac. Fish. Hokkaido Univ. 9: 29-36.
- Théel, H.J., 1876: Note sur l'*Elpidia*, genre nouveau du groupe des Holothurides. — Bih. K. svenska Vetensk. Akad. Handl. 4, 4: 1-7.
- 1877: Mémoire sur l'*Elpidia*, nouveau genre d'Holothurides. — K. svenska Vetensk. Akad. Handl. 14, 8: 1-30.
- 1879: Preliminary report on the Holothuridae of the Exploring voyage of H. M. S. "Challenger", I. — Bih. K. svenska Vetensk. Akad. Handl. 5, 19: 1-20.
- 1882: Report on the Holothurioidea, I. — Rep. scient. Results Voyage Challenger, Zool. 4, 13: 1-176.
- 1886a: Report on the Holothurioidea, II. — Ibid. 14, 39: 1-290.
- 1886b: Report on the Holothurioidea. Reports on the results of dredging... "Blake", 30. — Bull. Mus. comp. Zool. Harv. 13: 1-21.
- Thorson, G., 1946: Reproduction and larval development of Danish marine bottom invertebrates. — Meddr Kommn Danm. Fisk.- og Havunders., Ser. Plankton 4: 1-523.
- 1952: Zur jetzigen Lage der marinen Bodentier-Ökologie. — Zool. Anz., Suppl. 16: 276-327.
- Ursin, E., 1960: A quantitative investigation of the echinoderm fauna of the central North Sea. — Meddr Danm. Fisk.- og Havunders., N. S. 2, 24: 1-204.
- Ushakov, P. V., 1952: Investigations on the deep-sea fauna. — Priroda, Mosk. 1952, No. 6: 100-102. (In Russian).
- Vaney, C., 1906: Holothurides. — Expédition Antarctique Française (1903-1905). 30 pp.
- 1908: Les Holothurides de l'Expédition Antarctique Nationale Écossaise. — Trans. R. Soc. Edinb. 46: 405-441.
- 1914: Holothurides. — Deuxième Expédition Antarctique Française (1908-1910). 54 pp.
- Vasseur, E., 1952: Geographic variation in the Norwegian sea-urchins, *Strongylocentrotus droebachiensis* and *S. pallidus*. — Evolution, Lancaster, Pa. 6: 87-100.
- Verrill, A. E., 1884: Notice of a remarkable marine fauna occupying the outer banks off the southern coast of New England, No. 9. — Am. J. Sci. 28: 213-220.
- Verrill, A. E., 1885: Results of the explorations made by the steamer "Albatross" off the northern coast of the United States in 1883. — United States Commission of Fish and Fisheries, Part XI. Report of the Commissioner for 1883: 503-699.
- Vinogradova, N. G., 1959: The zoogeographical distribution of the deep-water bottom fauna in the abyssal zone of the ocean. — Deep Sea Res. 5: 205-208.
- 1962a: Vertical zonation in the distribution of deep-sea benthic fauna in the ocean. — Ibid. 8: 245-250.
- 1962b: Some problems of the study of deep-sea bottom fauna. — J. oceanogr. Soc. Japan, 20th Anniversary Volume: 724-741.
- Walsh, J. H. T., 1891: Natural history notes from H. M. Indian Marine Survey steamer "Investigator". No. 24. List of deep-sea holothurians, collected during seasons 1887-91, with descriptions of new species. — J. Asiatic Soc. Beng. 60: 197-204.
- Wolff, T., 1956a: Isopoda from depths exceeding 6000 meters. — Galathea Rep. 2: 85-157.
- 1956b: Crustacea Tanaidacea from depths exceeding 6000 meters. — Ibid. 2: 187-241.
- 1960: The hadal community, an introduction. — Deep Sea Res. 6: 95-124.
- 1961: Animal life from a single abyssal trawling. — Galathea Rep. 5: 129-162.
- 1962: The systematics and biology of bathyal and abyssal Isopoda Asellota. — Ibid. 6: 1-320.
- 1970: The concept of the hadal or ultra-abyssal fauna. — Deep Sea Res. 17: 983-1003.
- 1971: "Archimède" dive 7 to 4160 metres at Madeira: Observations and collecting results. — Vidensk. Meddr dansk naturh. Foren. 134: 127-147.
- Zenkevich, L. A., 1963: Biology of the seas of the USSR. — London. 955 pp.
- & Ya. Birshtein, 1956: Studies of the deep water fauna and related problems. — Deep Sea Res. 4: 54-64.
- — 1960: On the problem of the antiquity of the deep-sea fauna. — Ibid. 7: 10-23.
- Z. A. Filatova, G. M. Belyaev, T. S. Luk'yanova & I. A. Suetova, 1971: Quantitative distribution of zoobenthos in the world ocean. — Byull. mosk. Obshch. Ispyt. Prir. (Biol.) 76: 27-33. (In Russian, English summary).

VI. INDEX TO NAMES OF GENERA AND SPECIES

Main entries in *italics*. Generic names preceded by * refer to non-Elasipoda. Specific names in keys, diagrams of bathymetric distribution, and maps of geographic distribution are not included (see main entry to generic name).

* <i>Abyssocucumis</i> , 216	130, 150, 155-156, 159,	<i>Amperima</i> , 128, 129, 130,	186, 226, 235
<i>abyssorum</i> , 216, 234,	184, 187, 188, 190, 191,	158, 184, 186, 189, 192,	<i>tui</i> , 158
241	192, 213, 229	213, 229	<i>velacula</i> , 161, 185, 186,
* <i>Acanthotrochus</i> , 216	<i>ecalcareia</i> , 155, 156-157,	<i>furcata</i> , 159, 186, 226	215, 234
<i>mirabilis</i> , 216	185, 186, 187	<i>insignis</i> , 162	<i>Amphideima</i> , 15, 38
* <i>Acaudina</i> , 216	<i>lactea</i> , 156, 157	<i>naresi</i> , 159-161, 184,	<i>investigatoris</i> , 39, 43,
* <i>Achiridota</i> , 217	<i>monactinica</i> , 155,	185, 186, 192, 196, 204,	44, 45
<i>inermis</i> , 216	157-158, 187	242	* <i>Amphigymnas</i> , 215
<i>profunda</i> , 216	<i>paradoxa</i> , 156, 157	<i>robusta</i> , 161-162, 189	<i>Apodogaster</i> , 47, 75, 190,
<i>Achlyonice</i> , 127, 128, 129,	<i>tui</i> , 158, 187, 215	<i>rosea</i> , 158-159, 185,	192, 210, 215

- alcocki*, 75, 234
 **Astichopus*, 204
 multifidus, 204
Bathygone, 52
 papillatum, 67, 69, 70
 **Bathyplotes*, 186, 215, 216
 moseleyi, 234
 natans, 204, 234
 reptans, 234
Benthodytes, 10, 76-77,
 190, 191, 192, 194, 206,
 211, 225, 226
 abyssicola, 76
 assimilis, 106, 108
 browni, 76
 gigantea, 76
 glutinosa, 89
 gotoi, 77, 79, 80
 hystrix, 84, 85, 86, 87
 incerta, 76, 77-80, 87,
 191, 203
 janthina, 76, 80, 82, 89,
 93
 kerhervei, 102, 105
 lingua, 76, 80-82, 87,
 190, 191, 203, 214, 215
 mamillifera, 76
 papillifera, 89, 92, 93
 plana, 76, 84, 87-89,
 235
 recta, 99
 regularis, 77, 80
 salivosus, 75
 sanguinolenta, 76, 93,
 94-96, 190, 191, 194,
 203, 226
 selenkiana, 76
 sibogae, 76, 84-87, 203,
 220
 sordida, 76
 spuma, 76
 superba, 76, 89, 190
 typica, 76, 89-93, 187,
 191, 194, 202, 203, 204,
 205, 223, 227, 237, 238
 valdiviae, 76, 82-84
Benthogone, 47, 48, 52,
 190, 206, 210, 224
 abstrusa, 51-52, 220
 fragilis, 49-50, 191
 quadrilineata, 48, 49
 quatrolineata, 67
 rosea, 48-49, 50, 187,
 191, 202, 221
Benthophyces, 48, 50
 fragilis, 49, 50
 **Benthothuria*, 215
 **Capheira*, 215
 **Caudina*, 216
 chilensis, 189
 **Ceraplectana*, 216
 trachyderma, 216
 **Chiridota*, 216
 abyssicola, 216
Cryodora, 52
 spongiosa, 58
 **Cucumaria*, 189
 echinata, 189
Deima, 15, 16, 183, 190,
 191, 193, 205, 206, 210,
 222
 atlanticum, 18, 20, 22
 blakei, 17, 20, 21
 fastosum, 17, 20, 21,
 22, 23, 194
 mosaicum, 18, 20, 22, 23
 pacificum, 20, 24
 validum, 16-24, 190,
 191, 194, 198, 199, 215,
 223, 227, 228, 237, 241,
 245
 v. pacificum, 17, 22, 23,
 24
 v. validum, 17-23, 204
 **Dendrothuria*, 215
 **Ekmocucumis*, 216
 turqueti, 234
Ellipinion, 128, 129, 130,
 131, 162-163, 184, 186,
 189, 192, 213, 230
 bucephalum, 163-164
 delagei, 163, 215
 facetum, 166, 189
 galathea, 165, 192, 241
 kumai, 162, 166, 189,
 215
 molle, 165, 166, 169
 papillosum, 165-166,
 189
 solidum, 163, 164-165
Elpidia, 9, 10, 127, 128,
 130, 172-174, 184, 189,
 190, 192, 196, 205, 207,
 213, 227, 231, 232, 241,
 243
 adenensis, 172, 175, 232
 ambigua, 151
 antarctica, 174
 atakama, 172, 181, 182,
 232, 241, 242
 birsteini, 172, 181, 232,
 242, 243
 chilensis, 172, 175, 232
 glacialis, 8, 172, 173,
 175-181, 185, 186, 190,
 196, 204, 205, 215, 223,
 232, 234, 237, 242, 245
 g. glacialis, 175,
 176-178
 g. kermadecensis, 180
 g. kurilensis, 180-181,
 243
 g. solomonensis, 180,
 181
 g. sundensis, 178-180
 g. theeli, 174
 g. uschakovi, 180
 globosa, 167
 hanseni, 172, 181, 242,
 243
 h. hanseni, 181, 232
 h. idzubonensis, 181,
 232
 incerta, 143
 javanica, 178, 180
 kermadecensis, 180
 kurilensis, 180, 243
 longicirrata, 172, 181,
 232, 241, 243
 minutissima, 172, 175,
 232
 mollis, 165
 murrayi, 167
 nana, 170
 papillosa, 165
 purpurea, 151
 rigida, 136
 solomonensis, 180
 sundensis, 178
 theeli, 172, 174-175,
 185, 186, 232
 uschakovi, 180
 verrucosa, 147
 willemoesi, 144
Elpidiogone, 128, 129, 131
 dubia, 144
Eynpniastes, 127, 128
Euphronides, 99
 anchora, 102, 103, 104,
 105
 auriculata, 106, 108
 bifurcata, 102, 112
 cornuta, 106, 109, 110
 depressa, 106, 108, 109,
 110
 dyscrita, 116, 124
 kerhervei, 102, 105
 scotiae, 111
 talismani, 106, 108, 109,
 110
 tanneri, 106, 108, 109,
 110
 verrucosa, 112
 violacea, 106, 108, 109,
 110
 **Eupyrgus*, 216
 pacificus, 216
 scaber, 216
Euriplastes, 128, 132
 **Filithuria*, 215
 **Galatheathuria*, 185, 204
 aspera, 185, 204, 205
 **Gastrothuria*, 215
 **Gephyrothuria*, 216
 **Hadalothuria*, 195, 216
 wolffi, 242
 **Hedingia*, 216
 albicans, 216
 californica, 216
 fusiforme, 216
 ludwigi, 216
 planapertura, 216
 **Heteromolpadia*, 216
 marenzelleri, 216
 piki, 216
 tridens, 216
 **Holothuria*, 184
Ilyodaemon, 48, 52
 abstrusus, 48, 51, 190
 fimbriatus, 64, 67, 69
 ijimai, 67
 maculatus, 63
 miurense, 63, 65, 66
Irpa, 8, 127, 128, 130, 131,
 171, 186, 189, 192, 194,
 213, 230
 abyssicola, 171, 172,
 232
 ludwigi, 171, 172, 185,
 186, 189, 215, 235
 **Karenilla*, 215
Kolga, 8, 127, 128, 130,
 131, 170, 186, 187, 188,
 192, 194, 213, 230
 foliacea, 152
 furcata, 159
 hyalina, 170-171, 187,
 188, 205, 225, 232
 ludwigi, 172
 nana, 170, 171, 188
 obsoleta, 134
Laetmenoeus, 52
 scotoeides, 52, 61
Laetmogone, 47, 48, 52-
 53, 191, 210, 224, 225
 biserialis, 52, 53, 69,
 70-71, 191, 220, 238
 brongniarti, 58
 enisus, 64, 65
 fimbriata, 52, 53, 67-
 70, 71, 191, 202, 215,
 221, 228, 238
 ijimai, 52, 66, 67, 192,
 215
 interjacens, 47, 52, 53-
 54, 191, 220
 jourdaini, 58
 maculata, 52, 63-67,
 191, 202, 215, 228, 238
 neglecta, 70, 71
 parva, 67, 69, 70
 scotoeides, 52, 61-63,
 220
 selenkai, 67, 69, 70, 71
 spongiosa, 58
 theeli, 52, 57-58, 61,
 191, 215, 246
 violacea, 52, 57, 58-61,


- 185, 191, 201, 202, 215,
221, 234, 238
wyvillethomsoni, 52,
53, 54-57, 58, 61, 70,
71, 191, 215, 246
Laetmophasma, 72
fecundum, 72, 74
**Leptosynapta*, 189
inhaerens, 189
**Meseres*, 215
**Mesothuria*, 215
bifurcata, 234
candelabri, 241
intestinalis, 235
verrilli, 235
**Molpadia*, 216, 246
antarctica, 234
bathybia, 216
blakei, 216, 241, 246
granulata, 216, 246
musculus, 216, 235
parva, 216
**Myriotrochus*, 10, 210,
216, 217
bathybius, 216
bruuni, 242
giganteus, 216
kermadecensis, 242
kurilensis, 241
longissimus, 241
mitis, 242
theeli, 216
zenkevitchi, 242
Nectothuria, 99
translucida, 116, 124,
125
Oneirophanta, 15, 16, 24,
183, 191, 205, 206, 210,
215, 222
affinis, 32, 36
alternata, 25, 31
conservata, 36, 222
mutabilis, 24-36, 187,
190, 193, 194, 198, 214,
223, 226, 231, 232, 238,
241, 245
m. affinis, 32-36, 191,
199, 200, 201, 204, 222
m. mutabilis, 24-32,
199, 201, 222
setigera, 36-38, 47, 186,
194, 198, 222
Orphmurgus, 15, 16, 38,
191, 206, 210, 215, 220,
223
asper, 16, 38-39, 221
glaber, 16, 39-46, 183,
190, 194, 198, 220, 228,
238, 245
insignis, 16, 39, 43, 44,
45, 46, 47
invalidus, 39, 43, 45
protectus, 16, 47
rigidus, 39, 43, 45, 46
vitreus, 16, 46-47, 220
**Paelopatides*, 76, 204, 215
aspera, 185
**Paelpidia*, 208
norica, 208
Pannychia, 47, 72, 192,
210, 224
glutinosa, 80, 82
moseleyi, 72-75, 202,
220, 238
m. mollis, 72, 73
m. virgulifera, 72, 73,
74
multiradiata, 72, 74
pallida, 72, 74
**Paracaudina*, 216
Paelpidia, 127, 128, 129,
131, 190
anamesa, 132
cylindrica, 147
elongata, 147
**Paroriza*, 241
grevei, 241
Pelagothuria, 128
Peniagone, 127, 128, 129,
130, 131-134, 187, 188,
192, 194, 195, 205, 207,
212, 215, 228, 229
affinis, 139, 142-143,
187
atrox, 132
azorica, 138-142, 146,
187, 204, 206, 215, 242
bispiculata, 132
challengeri, 146
convexa, 135-136, 152,
187
diaphana, 152, 153-155,
187, 192, 205, 238
discrepans, 131
dubia, 144-145
ecalcareia, 131
elongata, 147, 148, 187
expansa, 152, 153
ferruginea, 151
foliacea, 152, 153
gracilis, 152, 155
horrifera, 132
humilis, 138
incerta, 139, 142,
143-144
incondita, 145
intermedia, 152-153
islandica, 150, 191
japonica, 147, 187, 215,
233
lacinora, 151
lugubris, 148, 151
mossmani, 134
mus, 132
naresi, 159
nybelini, 134
obscura, 132
obsoleta, 134-135, 136,
152
papillata, 145-146, 187
piriei, 132
porcella, 134
purpurea, 151-152, 153,
155, 186, 187, 205
rigida, 136-137, 138,
152
setosa, 148, 149
stabilis, 132
theeli, 146, 147, 185
vedeli, 138, 139
verrucosa, 147-148
vexillum, 132
vignoni, 146-147, 215,
233
vitrea, 143, 148-150,
187
willemoesi, 138, 139,
142, 144, 187
willtoni, 134
wyvillii, 131, 150-151
Periamma, 128, 158
furcata, 159
kumai, 166
ludwigi, 172
naresi, 159
roseum, 158
tetramerum, 159, 160
**Praeuephronides*, 209
multiperforata, 209
**Protankyra*, 217
abyssicola, 217
brychia, 217
**Pseudostichopus*, 215
mollis, 234
occultatus, 235
villosus, 242
**Pseudothuria*, 215
**Psolidium*, 216
disjunctum, 216
panamense, 216
**Psolus*, 216
phantapus, 189, 196
pourtalesii, 216
Psycheotrepes, 76, 96,
191, 192, 211, 227
exigua, 96, 98, 99, 191,
203
magna, 96, 97-99, 194,
241
recta, 96, 99
Psychrelpidia, 128, 129,
130, 131, 187, 190, 192,
194, 195, 206, 207, 212,
227
discrepans, 131
Psychropotes, 76, 99, 191,
192, 196, 204, 211, 226,
227
belyaevi, 100-102, 194,
203, 204, 235, 244
brucei, 116, 122, 123,
124, 125
buglossa, 116, 122
depressa, 106-111, 187,
191, 192, 194, 203, 204,
214, 215, 223, 234
dubiosa, 116, 124
fucata, 116, 122
grimaldii, 116, 122
kerhervei, 102, 105
laticauda, 116, 122,
123, 124, 125
longicauda, 99, 102,
115-126, 127, 182, 187,
190, 191, 194, 203, 205,
214, 223, 226, 232, 238,
241
loveni, 126-127, 191
minuta, 106
mirabilis, 102, 111-112,
115
rariipes, 116, 124
scotiae, 111
semperiana, 102-105,
106, 192, 194, 203
verrucosa, 102, 112-115,
194, 203
Rhipidothuria, 128, 129,
130, 131, 190, 192, 194,
212, 227
racowitzai, 131
Scotoanassa, 127, 128,
129, 131, 152
diaphana, 153
gracilis, 155
incerta, 132, 228
translucida, 153, 154
Scotodeima, 15, 38
parvispiculatum, 15
protectum, 47
setigerum, 36
vitreum, 46
Scotoplanes, 9, 128, 129,
130, 131, 166, 168, 184,
186, 189, 190, 192, 205,
213, 230, 234
albida, 162
angelicus, 165, 166
clarki, 166, 169-170,
184, 189, 196, 204, 234,
246
delagei, 162, 163
facetus, 166
galathea, 165
gilpibrowni, 155
globosa, 166, 167-169,
170, 184, 189, 196, 204,
215, 223, 232, 234,

242, 246
insignis, 162
mollis, 165
murrayi, 167, 168, 169
papillosa, 165
robusta, 161
theeli, 167, 168, 169
 **Sphaerothuria*, 216
 bitentaculata, 216
 **Staurocucumis*, 216
 liouvillei, 182, 208, 234

**Stichopus*, 186
 japonicus, 197
 regalis, 215, 235
 tremulus, 215
 **Synallactes*, 215, 216
 aenigma, 15, 216
 alexandri, 215
 anceps, 216
 carthagei, 234
 challengeri, 215
 chuni, 215
 crucifera, 216

discoidalis, 215
gilberti, 216
heteroculus, 215
horridus, 216
ishikawai, 215
mollis, 215
monoculus, 215
multivesiculatus, 215
nozawai, 215
profundus, 216
rigidus, 216
robertsoni, 216

triplax, 215
viridilimus, 215
 **Synaptula*, 189
 hydriformis, 189
Triconus, 99
 **Trochoderma*, 216
 elegans, 216
 **Trochostoma*, 246
 granulatum, 246
Tutela, 172
 echinata, 175



Additions and corrections

P. 76, right column, line 6 from below: Koehler & Vaney, 1905 Read Vaney, 1908.
 P. 127, left column, line 3 from below: Théel, 1879 Read Théel, 1882.
 P. 235, left column. With regard to the penetration of species into the Mediterranean deep sea the following information from E. Tortonese 1965 (Fauna d'Italia, 6) should be noted: (1) *Mesothuria verrilli* may not occur in the Mediterranean, the recorded specimens probably belonging to *M. intestinalis*. (2) *Molpadia musculus* penetrates to at least 820 m. (3) The molpadonian *Hedingia mediterranea* is known from the type specimen only (Tyrrhenian Sea: 800–1005 m). The species is likely to be identical with *H. albicans* (North Atlantic, Bay of Bengal, southwestern Pacific: 491–3200 m).