

REFERENCES

- Filatova, Z. A., M. N. Sokolova & R. Ya. Levenstein, 1968: Mollusc of the Cambro-Devonian Class Monoplacophora found in the Northern Pacific. — *Nature* **220**: 1114-1115.
- , N. G. Vinogradova & L. I. Moskalev, 1974: New finding of the ancient primitive mollusc *Neopilina* in the Atlantic part of the Antarctic. — *Ibid.* **249**: 657.
- Hedley, R. H. & K. M. Rudall, 1974: Extracellular silk fibres in *Stannophyllum* (Rhizopodea: Protozoa). — *Cell Tiss. Res.* **150**: 107-111.
- Lemche, H. 1957: A new living deep-sea mollusc of the Cambro-Devonian Class Monoplacophora. — *Nature* **179**: 413-416.
- Lemche, H. & K. G. Wingstrand, 1959: The anatomy of *Neopilina galathea* Lemche, 1957 (Mollusca Tryblidiacea). — *Galathea Rep.* **3**: 7-71.
- — 1960: Classe des monoplacophores. — *Traité de Zoologie* **5**, 2: 1787-1821.
- Lemche, H., B. Hansen, F. Jensenius Madsen, O. S. Tendal & T. Wolff, 1976: Hadal life as analyzed from photographs. — *Vidensk. Meddr dansk naturh. Foren.* **139**: 263-336.
- Lowenstam, H. A., 1978: Recovery, behaviour and evolutionary implications of live Monoplacophora. — *Nature* **273**: 231-232.
- Marshall, N. B., 1979: *Developments in Deep-Sea Biology*. — Blandford Press, 566 pp.
- McLean, J. H., 1979: A new monoplacophoran limpet from the continental shelf off Southern California. — *Contrib. Sci. nat. Hist. Mus. Los Angeles No.* 307: 1-19.
- Menzies, R. J., M. Ewing, J. L. Worzel & A. H. Clarke, 1959: Ecology of the recent Monoplacophora. — *Oikos* **10**: 168-182.
- , R. Y. George & G. T. Rowe, 1973: *Abyssal Environment and Ecology of the World Oceans*. — John Wiley & Sons, New York, 488 pp.
- Moskalev, L. I., Ya. I. Starobogatov & Z. A. Filatova, 1983: New data on the abyssal Monoplacophora from the Pacific and South Atlantic Oceans. — *Zool. Zh.* **62**: 981-996 (In Russian).
- Rokop, F. J., 1972: A new species of monoplacophoran from the abyssal North Pacific. — *Veliger* **15**: 91-95.
- Tendal, O. S., 1972: A monograph of the Xenophyophoria (Rhizopodea, Protozoa). — *Galathea Rep.* **12**: 7-99.
- 1979: Aspects of the biology of Komokiacea and Xenophyophoria. — *Sarsia* **64**: 13-17.
- Wingstrand, K. G., 1985: On the anatomy and relationships of recent Monoplacophora (Mollusca). — *Galathea Rep.* **16**: 7-94.
- Wolff, T., 1961: Animal life from a single abyssal trawling. — *Ibid.* **5**: 129-162.
- Younge, C. M., 1957: Reflexions on the monoplacophoran, *Neopilina galathea* Lemche. — *Nature* **179**: 672-673.
- 1960: General characters of Mollusca. — In: *Treatise on Invertebrate Paleontology Part 1, Mollusca* 1: 3-36.

PLATE 13

- A. Part of *Stannophyllum zonarium* with numerous scrape marks. Bar represents 1 cm. Arrows point to some of the scrape marks.
- B. Cross section of first intestinal coil of *Neopilina galathea*. Spec. III. Bar represents 100 μ m. s, stercomare fragment. g, granellare fragment.
- C. Section of stercomare fragment from first intestinal coil. Numerous xenophyophore fecal pellets (stercomata) surrounded by membrane (clearly seen to the right). Bar represents 100 μ m.
- D. Section of granellare from first intestinal coil. Xenophyophore plasma containing numerous barite crystals (granellae) and surrounded by membrane. Bar represents 20 μ m.
- E. Section of granellare from fourth intestinal coil. Granellae in nearly empty granellare membrane. Bar represents 50 μ m.

